



Evagio[®] Forte

syngenta.

GROUP 21 40 FUNGICIDES



Product registration number: PCS No. 06838
UF: 9PQ7-F0SV-000J-XCTR

A suspension concentrate fungicide containing
100 g/L of amisulbrom, 200 g/L of mandipropamid.

For the control of late blight (*Phytophthora infestans*)
in potatoes.

Authorisation holder

Syngenta UK Limited,
Jealott's Hill International Research Centre, Bracknell,
Berkshire, RG42 6EY
Tel +44 (0) 1223 883400

Marketing Company

Syngenta Ireland Limited
Block 6, Cleaboy Business Park, Old Kilmeaden Road,
Waterford
Tel: (051) 377203

In case of toxic or transport emergency ring

+44 (0) 1484 538444 (24hr)

STORE IN A COOL, DRY PLACE.
PROTECT FROM FROST.

© Syngenta 2026



FOR PROFESSIONAL USE ONLY

To avoid risks to human health and the environment comply
with the instructions for use.

EVAGIO[®] Forte

A suspension concentrate containing 100 g/L
of amisulbrom, 200 g/L of mandipropamid.

Warning

Causes serious eye irritation.

Suspected of causing cancer.

**Very toxic to aquatic life with long
lasting effects.**

Keep out of reach of children.

Obtain special instructions before use.

Avoid release to the environment.

Wear protective gloves/protective clothing/
eye protection/face protection/hearing protection.

If exposed or concerned: get medical advice/attention.

If eye irritation persists: get medical advice/attention.


Collect spillage.

Dispose of contents/container to a licensed hazardous-
waste disposal contractor or collection site except for triple
rinsed empty containers which can be disposed of as non-
hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

PCS 06838 UF: 9PQ7-F0SV-000J-XCTR

L1144885 IREL/10D PPE 4243496



5 litres

Product names marked © or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (L/ha)	Maximum total dose (L/ha)	Maximum number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Potatoes and seed potatoes	0.6	1.8	3	6	7 days before harvest

READ LABEL BEFORE USE: USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE: FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS**(a) Operator protection**

WEAR SUITABLE EYE PROTECTION AND PROTECTIVE CLOTHING (gloves, coveralls) when handling the product.

KEEP OUT OF THE REACH OF CHILDREN.

Do not eat, drink or smoke when using the product.

WASH HANDS AND EXPOSED SKIN before meals and after work.

(b) Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

(c) Storage and Disposal

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

This leaflet is part of the approved Product Label.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

EVAGIO® FORTE can be used on all varieties of potatoes including seed potatoes.

PROPERTIES OF EVAGIO FORTE

EVAGIO FORTE is a foliar fungicide for the preventative control of late blight (*Phytophthora infestans*) in potatoes and is highly active against spore germination. Uptake into the leaf tissues assures good translaminar and local systemic activity and inhibits mycelial growth during the incubation period.

DISEASE CONTROLLED

Control of late blight (*Phytophthora infestans*) in potatoes.

BACKGROUND TO POTATO BLIGHT CONTROL

Late blight (*Phytophthora infestans*) is potentially a devastating disease of potatoes. In commercial production, a season long disease prevention policy is essential. First of all ensure that other control methods are being satisfactorily applied:

1. Prevent re-growth on potato dumps.
2. Destroy all groundkeepers.
3. Plant disease free seed.
4. Use generous soil ridges to protect tubers.

Integrate this approach with a fungicide programme:

Early Crops

In first and second early potato crops, particularly those grown in the same locality as main crop potatoes, an adequate and full blight protection programme should be applied right up to harvesting or haulm desiccation. This will protect the early crop while helping to reduce disease risk to later crops.s.

Maincrops

Disease prevention programmes require regular and season long fungicide use to limit foliar blight development. However, as an effective fungicide programme will preserve leaf area there may be more risk of infecting tubers at harvest, particularly during “heavy” blight years. Completion of the control programme should therefore include a complete haulm desiccant. Lifting of the crop should not take place for at least 10 days after COMPLETE KILL of the haulm. Crops intended for storage should not be lifted while there is any green tissue AT ALL on the leaves or stem bases.

Blight Risk Assessment

The risk of disease is affected by **weather conditions** (during the crop life) and **crop location**:

Weather Conditions

Spread of disease occurs under warm, humid conditions. Preferably use a reliable decision support system to determine what frequency of fungicide treatment is appropriate and fungicide type required.

Crop Location

Locations with the highest probability of blight problems are:

1. Areas of the country where extensive main crop or early production takes place
2. Areas where climatic conditions that encourage disease development occur on a frequent basis

TIMING

EVAGIO FORTE is a protectant and curative fungicide so following good agricultural practice the programme should start BEFORE blight enters the crop. Commence spraying at the first blight warning or when local weather conditions are favourable for the disease. However applications 1 day after an infection event has occurred have shown to give good levels of control.

Intervals between applications of EVAGIO FORTE should be reduced as blight risk increases, so that protection of the crop can be maintained. Applications of EVAGIO FORTE should be made at 7-10 day intervals depending on disease pressure. As disease pressure and the risk of late blight infection increase, the interval should be shortened.

Rates of Use

Apply EVAGIO FORTE at 0.6 kg product per hectare. Up to three applications may be made per crop between growth stages 51 and 89. Allow a minimum of 6 days between applications. Applications of EVAGIO FORTE can be made up to 7 days before harvest.

MIXING AND SPRAYING

Spray Volume

Apply EVAGIO FORTE in a recommended 100-400 litres of water per hectare.

Spray Nozzles

A medium quality spray* is preferred for application of EVAGIO FORTE (* as defined by The British Crop Protection Council). A spray pressure of 2 - 3 bar is recommended.

Mixing

Make sure the sprayer is clean and set to give an even application at the correct volume. Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of EVAGIO FORTE and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying and any stoppages.

Thoroughly wash all spray equipment with water immediately after use. Drain the system completely and rinse spray tank, boom and nozzles three times with clean water until the foam and all traces of product have been removed. Do not leave the spray liquid in the sprayer for long periods (such as during meal breaks or overnight).

RESISTANCE MANAGEMENT

EVAGIO FORTE contains mandipropamid, a CAA fungicide (FRAC code no. 40) and amisulbrom, a Qi-site inhibitor (Qil) fungicide (FRAC code no. 21). To minimise the risk of resistance development in the pathogen population the following guidelines for blight fungicides (based on an average number of 12 fungicide applications/season for blight control) should be followed:

1. Where possible, use an alternating strategy using fungicides from different mode of action groups.
2. Where CAA fungicides are applied as a mixture (co-formulated or as a tank mix) up to six applications (or max. of 50% of the total number of applications) may be made per crop or season.
3. No more than 3 applications of any CAA fungicide should be made consecutively.
4. Further information on suitable tank mix products and resistance management strategies is available from FRAG and BCPC websites.

COMPANY ADVISORY INFORMATION

As part of our Product Stewardship policy, Syngenta recommends the following precautions should also be observed:

Wear appropriate clothing – cotton overall and rubber gloves, when handling the concentrate.

Since the occurrence of resistance cannot be forecast, neither Syngenta UK Limited nor its distributors can accept responsibility for any loss or damage to crops caused by the failure of EVAGIO FORTE to control resistant strains.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (L/ha)	Maximum total dose (L/ha)	Maximum number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Potatoes and seed potatoes	0.6	1.8	3	6	7 days before harvest

READ LABEL BEFORE USE: USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE: FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS**(a) Operator protection**

WEAR SUITABLE EYE PROTECTION AND PROTECTIVE CLOTHING (gloves, coveralls) when handling the product.

KEEP OUT OF THE REACH OF CHILDREN.

Do not eat, drink or smoke when using the product.

WASH HANDS AND EXPOSED SKIN before meals and after work.

(b) Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

(c) Storage and Disposal

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.